arXiv:0706.0572v2 [hep-th] 21 Jun 2007

USTC-ICTS-07-09

Is Noncommutative Eternal Inflation Possible?

Yi-Fu Cai[1]*, Yi Wang[2,3]†

[1] Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, P. R. China

[2] Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, P. R. China

[3] The Interdisciplinary Center for Theoretical Study of China (USTC),
Hefei, Anhui 230027, P. R. China

Abstract

We investigate the condition for eternal inflation to take place in the noncommutative spacetime. We find that the possibility for eternal inflation's happening is greatly suppressed in this case. If eternal inflation can not happen in the low energy region where the noncommutativity is very weak (the UV region), it will never happen along the whole inflationary history. Based on these conclusions, we argue that an initial condition for eternal inflation is available from the property of spacetime noncommutativity.

1 Introduction

Inflation has been widely considered as a remarkably successful theory in explaining many problems in very early universe, such as the flatness, the horizon and the monopole problem [1, 2, 3, 4]. It predicts that the quantum fluctuations of the inflaton field were generated to form today's large scale structure[5, 6, 7, 8, 9], and

---

*caiyf@mail.ihep.ac.cn

†wangyi@itp.ac.cn

these fluctuations fit with current observations of cosmic microwave background very well[10, 11].

It is a common viewpoint that eternal inflation[12, 13, 14] can take place in a variety of inflation models. Especially, with the monomial chaotic inflaton potential, inflation generally becomes eternal in the high energy regime. Eternal inflation is not only important in concept, but also provides a possible realization for the string theory landscape[15, 16, 17, 18]. There have been an increasing number of papers investigating whether we can measure eternal inflation and calculate probabilities in the multiverse, see references[19, 20, 21, 22, 23, 24].

Usually, eternal inflation happens when the energy scale of the universe is extremely high. Thus we would like to take into consideration on more fundamental theories in logic, namely, the string theory. In this paper, we focus on a universal property of string theory, i.e., noncommutativity of spacetime. By considering the effects of spacetime noncommutativity, we study the condition for eternal inflation and its implications. As required by stringy spacetime uncertainty relation[25, 26, 27], the physical time t_p and the physical length x_p should satisfy

$$\Delta t_p \Delta x_p \geq l_s^2 \ , \tag{1}$$

where l_s is a length scale given by the string theory.

There have been a lot of attempts to apply this uncertainty relation into inflationary cosmology, dubbed noncommutative inflation, see Ref. [31, 32, 33]. More detailed works have been investigated in a number of literature[34, 35, 36, 37, 38, 39, 40]. Here we briefly review the conception of noncommutative inflation proposed by Brandenberger and Ho[33], and then examine whether eternal inflation takes place with such noncommutativity. Note that here we have slightly generalized the discussion in [33] from power law inflation into a generally quasi-exponential inflationary scenario.

In order to introduce the noncommutativity into the 4-dimensional Friedmann-Robertson-Walker universe, we would like to define another time coordinate τ,

$$ds^2 = dt^2 - a^2(t)d\vec{x}^2 = a^{-2}(\tau)d\tau^2 - a^2(\tau)d\vec{x}^2 \ , \tag{2}$$

where a is the scale factor and we have assumed a spatially flat universe($K = 0$). For the quasi-exponential expansion where we have imposed the usual slow roll approximation self-consistently, we have

$$d\tau = adt \simeq a_0 e^{Ht} dt \ , \quad a \simeq H\tau \ , \tag{3}$$

where $H = \dot{a}/a$ is the Hubble parameter with the dot represents the derivative with respect to the cosmic time t and a_0 is an arbitrary parameter for the scale factor at a fixed time. Then the spacetime uncertainty relation takes the form

$$\Delta\tau\Delta x \geq l_s^2 \ , \tag{4}$$

where x is the comoving radial coordinate. This can be realized by the commutation relation of spacetime

$$[\tau, x]_* = il_s^2 \ , \tag{5}$$

where the $*$-product is defined as

$$(f * g)(x,\tau) = \exp\left(-\frac{i}{2} l_s^2 \left(\partial_x \partial_{\tau'} - \partial_\tau \partial_y\right)\right) f(x,\tau) g(y,\tau')\Big|_{y=x,\tau'=\tau} \ . \tag{6}$$

Based on these considerations, for the fluctuation of the inflaton field $\delta_q \varphi_k$, we can derive that the canonical normalized perturbation variable $u_k \simeq a\delta_q \varphi_k$ satisfies the equation of motion

$$u_k{}'' + \left(k^2 - \frac{z_k{}''}{z_k}\right) u_k = 0 \ , \tag{7}$$

where the prime denotes the derivative with respect to $\tilde{\eta}$ defined as $d\tilde{\eta} \equiv a_{\text{eff}}^{-2} d\tau$. The parameter a_{eff} is an effective scale factor appeared in the dispersion relation between a mode k and its energy defined with respect to τ,

$$a_{\text{eff}}^2 \equiv \left(\frac{\beta_k^+}{\beta_k^-}\right)^{\frac{1}{2}} = \frac{k}{E_k} \ , \quad \beta_k^{\pm}(\tau) \equiv \frac{1}{2}\left[a^{\pm 2}(\tau - l_s^2 k) + a^{\pm 2}(\tau + l_s^2 k)\right] \ , \tag{8}$$

and z_k is defined as

$$z_k^2 \equiv (\beta_k^- \beta_k^+)^{\frac{1}{2}} z^2 \ , \quad z \equiv a\frac{\dot{\varphi}}{H} \ . \tag{9}$$

As the scale factor is expanding nearly exponentially, when $l_s^2 k$ is not too small compared with τ we can take the approximate form of $\beta_k^{\pm}$ as

$$\beta_k^+ \simeq \tfrac{1}{2} a^2(\tau + l_s^2 k) \simeq \tfrac{1}{2}\left[H \cdot (\tau + l_s^2 k)\right]^2 \ , \tag{10}$$

$$\beta_k^- \simeq \tfrac{1}{2} a^{-2}(\tau - l_s^2 k) \simeq \tfrac{1}{2}\left[H \cdot (\tau - l_s^2 k)\right]^{-2} \ . \tag{11}$$

Moreover, because of the relations $\Delta x \sim 1/k, \Delta\tau \sim 1/E_k$ and the spacetime uncertainty (4), there is an initial time τ_k for the perturbations to be generated,

$$a_{\text{eff}}(\tau) \geq a_{\text{eff}}(\tau_k) = l_s k \ , \tag{12}$$

and the fluctuations are not allowed to exist before τ_k.

Due to this initial time for the fluctuations, the quantum fluctuations of inflaton in noncommutative environment can be generated inside or outside the Hubble radius. These fluctuations are called UV modes and IR modes respectively. In the UV mode limit, the effects of spacetime noncommutativity becomes very weak and hence the evolution of the fluctuations is similar with the commutative case. Therefore the spectrum of UV modes is roughly the same as in the commutative case. While in the IR region, noncommutativity dominates the behavior of the perturbations, and some familiar pictures will be totally modified. In this paper, we shall use the analysis of noncommutativity mentioned above (especially the IR region) to study eternal inflation.

This paper is organized as follows. In Section 2, we calculate the fluctuations generated outside the horizon and provide the constraint for eternal inflation to exist. In Section 3, we make a conclusion and discuss some other possibilities.

2 Fluctuations generated outside the horizon

In this section, we investigate the quantum fluctuations generated in noncommutative inflationary era and discuss the condition of eternal inflation. Since in the UV case the noncommutative property only contribute a few corrections on the usual perturbation theory[33, 34], the condition for the eternal inflation to happen is basically the same as the commutative one. Therefore, we focus our consideration on the IR case that the quantum fluctuations are generated outside the horizon. At the time the fluctuations start to be generated, the effective scale factor is

$$a_{\text{eff}}(\tau_k) = l_s k. \tag{13}$$

Making use of Eq. (10) and (11), and the fact that the Hubble scale is larger than the noncommutativity scale in this case, we get the initial time τ_k and the initial scale factor a_k as

$$\tau_k = \sqrt{l_s^4 k^2 \left(1 + \frac{1}{l_s^2 H^2}\right)} \simeq l_s^2 k, \quad a_k \simeq H l_s^2 k \tag{14}$$

Since the IR modes are generated outside the horizon, it is required that $k < a_k H$. By using the second relation in Eq. (14), we can see that

$$H > l_s^{-1} \tag{15}$$

in the IR case. This verifies the physical picture that H should be larger than the noncommutativity scale.

Now we calculate the quantum fluctuation in the momentum space $\delta_q \varphi_k$. $\delta_q \varphi_k$ is linked to the canonical perturbation u_k by $u_k \simeq a \delta_q \varphi_k$, and when the perturbation begins to be generated the initial u_k was canonically normalized as $u_k \simeq \frac{1}{\sqrt{2k}}$. Consequently, when $\delta_q \varphi_k$ is born its amplitude can be given by

$$\delta_q \varphi_k \simeq \frac{1}{\sqrt{2k}} \frac{1}{H l_s^2 k} \ . \tag{16}$$

After that, the fluctuations outside the horizon are nearly frozen. By transforming to the coordinate space, we obtain the relation

$$\left\langle \delta_q \varphi^2 \right\rangle = \int_{k=aH}^{k=(e\times a)H} \frac{dk}{k} \frac{k^3}{2\pi^2} \delta_q \varphi_k \delta_q \varphi_{-k} \simeq \left(\frac{1}{2\pi} \frac{1}{H l_s^2}\right)^2 \tag{17}$$

during one Hubble time, and correspondingly the IR quantum fluctuation $\delta_q \varphi$ per e-fold in the noncommutative spacetime is given by

$$\delta_q \varphi \simeq \frac{1}{2\pi} \frac{1}{H l_s^2} \ . \tag{18}$$

Note that this result is strongly different from the commutative one $\delta_q \varphi \simeq H/2\pi$. Due to this, the physics of eternal inflation is greatly modified in the noncommutative case and the condition for eternal inflation to happen needs to be reconsidered.

As usual, the classical motion of the inflaton during one Hubble time takes the form

$$\delta_c \varphi \simeq \dot{\varphi} H^{-1} \sim \frac{V_\varphi}{H^2} \ , \tag{19}$$

where V_φ denotes $dV(\varphi)/d\varphi$. And as usual, the condition that inflation becomes eternal is roughly $\delta_q \varphi > \delta_c \varphi$, or

$$H > V_\varphi l_s^2 \ . \tag{20}$$

Compared with the eternal inflation condition in the commutative case $H^3 > V_\varphi$, the noncommutative eternal inflation is more unlikely to happen. It is because the quantum fluctuation is generally smaller due to the suppression by the Hubble parameter. In the following, we shall consider two explicit examples with the chaotic potentials.

Firstly, let us consider the model $V = \frac{1}{2}m^2\varphi^2$. From the condition (15) we can see that the inflation is in the IR region when $\varphi > \frac{M_p}{l_s m}$ while in the UV region when $\varphi < \frac{M_p}{l_s m}$. Consequently, the condition that inflation has become eternal in the UV region is

$$\varphi_{\rm IR} \equiv \frac{M_p}{l_s m} > \sqrt{\frac{M_p^3}{m}}, \quad m < \frac{1}{M_p l_s^2} \ . \tag{21}$$

Interestingly, we note that from (20) the eternal inflation will nicely continue in the IR region if

$$m < \frac{1}{M_p l_s^2}. \tag{22}$$

To substitute the well-known value $m \sim 10^{-6}M_p$[41] into (22), we obtain the result that eternal inflation is allowed to happen only if $M_s \simeq l_s^{-1} > 10^{-3}M_p$.

So we conclude that if inflation has become eternal before entering the IR region, it can continue to be eternal in the IR region. But the amplitude of fluctuation can not grow to larger values because $\delta_q\varphi$ is now suppressed by a factor of $1/H$. This is greatly different from the commutative eternal inflation and may constrain the initial condition space for the eternal inflation. On the other hand, if inflation has not become eternal before entering the IR region, then the inequality (22) does not hold. In this case inflation will never become eternal because the quantum fluctuation can not be large enough.

As a second example, consider $V = \lambda M_p^{4-p}\varphi^p$ with $p > 2$. Then similarly, the condition for entering the IR region $H > 1/l_s$ requires

$$\varphi > \lambda^{-\frac{1}{p}} l_s^{-\frac{2}{p}} M_p^{\frac{p-2}{p}} \ . \tag{23}$$

From the inequality (20), we obtain the condition for inflation to be eternal in the IR region as follow,

$$\varphi < p^{-\frac{2}{p-2}} \lambda^{-\frac{1}{p-2}} l_s^{-\frac{4}{p-2}} M_p^{\frac{p-6}{p-2}} \tag{24}$$

In order that (23) and (24) has overlap and inflation can be eternal in the IR region,



Figure 1: Evolution of the chaotic inflation field with $V(\varphi) = \frac{1}{2}m^2\varphi^2$ in the noncommutative spacetime. Here we have assumed $M_s \simeq l_s^{-1}$ to be large enough so that eternal inflation can take place in the UV region. In this case, inflation is eternal in part of the noncommutative UV region and the whole IR region.

we need

$$\lambda < p^{-p}\left(\frac{1}{l_s M_p}\right)^{p+2} \quad . \tag{25}$$

If (25) is satisfied, the noncommutative eternal inflation is allowed to exist in the IR region, and there is an upper bound for the energy density of eternal inflation which arises from (24). Since the inflaton field can not climb higher than the bound (24) with large probability, this provides a possible initial condition for eternal inflation.

Now let us examine whether there is eternal inflation at all in the $p > 2$ model above. For the eternal inflation to take place in the UV region, we again obtain the relation

$$\lambda < p^{-p}\left(\frac{1}{l_s M_p}\right)^{p+2} \quad , \tag{26}$$

which is the same as the IR region eternal condition (25).

So we get the similar conclusion with the $\frac{1}{2}m^2\varphi^2$ ($p = 2$) model that if eternal inflation takes place in the UV region, it is allowed to extend into the IR region. On the other hand, if inflation can not be eternal in the UV region, unfortunately there will not be any eternal inflation in the whole noncommutative inflationary history. Besides, there is an interesting difference between $p > 2$ and $p = 2$ cases that, in the $p > 2$ case noncommutative eternal inflation requires an upper bound on the inflaton while this bound does not exist in the $p = 2$ case.

For example, in the $\lambda\varphi^4$ model, if we expect eternal inflation to take place, then we need the noncommutative scale $l_s^{-1} > \lambda^{\frac{1}{6}} M_p$. To apply the data $\lambda \sim 10^{-14}$, we obtain that this scale is around $10^{-2}M_p$. If l_s^{-1} is just above this scale, the upper bound (24) on φ is around $10^3 M_p$ and the corresponding energy density is about $10^{-2}M_p^4$. Consequently, we conclude that the limitation from noncommutativity can be much stronger than that from the Planck density. Such a string scale can be realized easily if we do certain compactification on a large manifold, which should be common and have a large prior probability in the string landscape.

On the other hand, such a scale $l_s^{-1} \sim 10^{-2}M_p$ or $l_s^{-1} \sim 10^{-3}M_p$ is considerably higher than the scale for the final 60 e-folds inflation. So if we can observe the noncommutativity in future CMB experiments[39], then eternal inflation is not expected to have happened.



Figure 2: Evolution of the chaotic inflation field with $V(\varphi) = \lambda\varphi^4$ in the noncommutative spacetime. Here we have assumed $M_s \simeq l_s^{-1}$ to be large enough so that eternal inflation can take place in the UV region. In this case, inflation is eternal in part of the noncommutative UV region and part of the IR region. This figure is different from Fig. 1 that eternal inflation can not happen in the green part of IR region where the energy scale is extremely high.

3 Conclusion and discussions

Spacetime noncommutativity, predicted by string theory, has become a fundamental principle and been studied in a number of literature (see e.g. [28, 29, 30]). This principle brings new physics when applied into inflation theory. In this paper, we have seen that for the chaotic inflationary potential, the scenario of eternal inflation in noncommutative spacetime is remarkably different from the usual one. If inflation do not become eternal in the UV region, then it can never have happened.

We have also discussed that if eternal inflation happens both in the UV region and the IR region, an initial condition for eternal inflation can be provided or constrained. From the derivation in this paper, we can see that $\delta_q\varphi$ becomes smaller and smaller along the potential when eternal inflation enters IR region. Therefore, for the $p = 2$ model the initial condition space is greatly reduced; while for the $p > 2$ model there is an upper bound for φ explicitly. Eternal inflation can not climb into higher energy regions than this bound. This provides an initial condition for eternal inflation. As is discussed in [20, 23], initial conditions may be essential for predictions in the multiverse. The initial condition discussed in this paper provides a possible solution for the initial condition problem and can be used in calculating the eternal inflationary probabilities.

In the derivation made in Section 2, we have used the standard method according to a number of calculations for noncommutative and non-eternal inflation in the literature. However, generally there are several possibilities which may change the results we obtain in this paper.

The simplest possibility is that there is no spacetime noncommutativity at all or the scale of noncommutativity is of the same order as the Planck scale. Therefore, the noncommutativity will not alter the picture of the usual eternal inflation.

As another possibility, the noncommutative field theory may be not precise enough to describe the generation of the quantum fluctuations. This saturation is somewhat similar to that inflationary fluctuations described by the common quantum field theory suffers from the transPlanckian problems[42]. To see this problem in the noncommutative field theory, we note that we have used the relation $\Delta x \sim 1/k$ in the calculations of noncommutative inflation. However, this relation does not hold in theories with certain UV-IR relations while UV-IR relations arise commonly together with noncommutativity. So there is possibility that the perturbations are generated even if $k > a_{\mathrm{eff}}/l_s$, but it is still an open issue for us to fully understand the physics in this region.

Finally, the background geometry may be affected considerably by noncommutativity in the IR region. However, up to now this case has not been carefully studied even in the non-eternal inflationary regime.

Acknowledgments

We thank Bin Chen, Miao Li, Yun-Song Piao and Xinmin Zhang for helpful discussions. This work is supported in part by National Natural Science Foundation of China under Grant Nos. 90303004, 10533010, 19925523 and 10405029, and by the Chinese Academy of Science under Grant No. KJCX3-SYW-N2. The author Y.W. acknowledgments grants of NSFC.

References

[1] A. H. Guth, Phys. Rev. **D23**, 347 (1981).

[2] A. D. Linde, Phys. Lett. **B108**, 389 (1982).

[3] A. Albrecht, and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).

[4] For earlier attemps on an inflationary model, see A. A. Starobinsky, JETP Lett. **30**, 682 (1979) [Pisma Zh. Eksp. Teor. Fiz. **30** (1979) 719] ; A. A. Starobinsky, Phys. Lett. **B91**, 99 (1980).

[5] V. Mukhanov, and G. Chibisov, JETP **33**, 549 (1981).

[6] A. H. Guth, and S.-Y. Pi, Phys. Rev. Lett. **49**, 1110 (1982).

[7] S. W. Hawking, Phys. Lett. **B115**, 295 (1982).

[8] A. A. Starobinsky, Phys. Lett. **B117**, 175 (1982).

[9] J. M. Bardeen, P. J. Steinhardt, and M. S. Turner, Phys. Rev. **D28**, 679 (1983).

[10] A. D. Miller *et al.*, Astrophys. J. **524**, L1 (1999), astro-ph/9906421; P. de Bernardis *et al.*, Nature **404**, 955 (2000), astro-ph/0004404; S. Hanany *et al.*, Astrophys. J. **524**, L5 (2000), astro-ph/0005123; N. W. Halverson *et al.*, Astrophys. J. **568**, 38 (2002), astro-ph/0104489; B. S. Mason *et al.*, Astrophys. J. **591**, 540 (2003), astro-ph/0205384; A. Benoit *et al.*, Astro. Astrophys. **399**, L25 (2003), astro-ph/0210306; J. H. Goldstein *et al.*, Astrophys. J. **599**, 773 (2003), astro-ph/0212517;

[11] D. N. Spergel *et al.*, Astrophys. J. Suppl. **148**, 175 (2003), astro-ph/0302209; D. N. Spergel *et al.*, astro-ph/0603449.

[12] P. J. Steinhardt, **The Very Early Universe** Proceedings, pp.251, (1982).

[13] A. Vilenkin, Phys. Rev. **D27**, 2848 (1983).

[14] A. D. Linde, Mod. Phys. Lett. **A1**, 81 (1986); A. D. Linde, Phys. Lett. **B175**, 395 (1986); A. S. Goncharov, A. D. Linde, and V. F. Mukhanov, Int. J. Mod. Phys. **A2**, 561 (1987); A. D. Linde, D. A. Linde and A. Mezhlumian, Phys. Lett. **B345**, 203 (1995)

[15] R. Bousso and J. Polchinski, JHEP **0006**, 006 (2000) [arXiv:hep-th/0004134].

[16] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, Phys. Rev. **D68**, 046005 (2003) [arXiv:hep-th/0301240].

[17] L. Susskind, arXiv:hep-th/0302219.

[18] M. R. Douglas, JHEP **0305**, 046 (2003) [arXiv:hep-th/0303194]; F. Denef and M. R. Douglas, JHEP **0405**, 072 (2004) [arXiv:hep-th/0404116]; M. R. Douglas and S. Kachru, arXiv:hep-th/0610102; F. Denef, M. R. Douglas and S. Kachru, arXiv:hep-th/0701050.

[19] J. Garriga, D. Schwartz-Perlov, A. Vilenkin and S. Winitzki, JCAP **0601**, 017 (2006) [arXiv:hep-th/0509184]; D. Schwartz-Perlov and A. Vilenkin, JCAP **0606**, 010 (2006) [arXiv:hep-th/0601162]; A. Vilenkin,

arXiv:hep-th/0602264; V. Vanchurin and A. Vilenkin, Phys. Rev. **D74**, 043520 (2006) [arXiv:hep-th/0605015]; A. Vilenkin, arXiv:hep-th/0609193.

[20] R. Bousso, Phys. Rev. Lett. **97**, 191302 (2006) [arXiv:hep-th/0605263]; R. Bousso, B. Freivogel and I. S. Yang, Phys. Rev. **D74**, 103516 (2006) [arXiv:hep-th/0606114].

[21] G. W. Gibbons and N. Turok, arXiv:hep-th/0609095.

[22] A. Linde, JCAP **0701**, 022 (2007) [arXiv:hep-th/0611043]; A. Linde, arXiv:0705.1160 [hep-th].

[23] D. Podolsky and K. Enqvist, arXiv:0704.0144 [hep-th].

[24] M. Li and Y. Wang, arXiv:0704.1026 [hep-th].

[25] M. Li and T. Yoneya, Phys. Rev. Lett. **78**, 1219 (1997) [arXiv:hep-th/9611072]; T. Yoneya, Prog. Theor. Phys. **103**, 1081 (2000) [arXiv:hep-th/0004074].

[26] A. Connes, M. Douglas, and A. Schwarz, JHEP **9802**, 003 (1998), hep-th/9711162.

[27] N. Seiberg, and E. Witten, JHEP **9909**, 032 (1999), hep-th/9908142.

[28] D. Bigatti, and L. Susskind, Phys. Rev. **D62**, 066004 (2000), hep-th/9908056; N. Seiberg, L. Susskind, and N. Toumbas, JHEP **0006**, 021 (2000), hep-th/0005040.

[29] A. Y. Alekseev, A. Recknagel, and V. Schomerus, JHEP **0005**, 010 (2000), hep-th/0003187.

[30] C.-S. Chu, and P.-M. Ho, Nucl. Phys. **B636**, 141 (2002), hep-th/0203186.

[31] C. S. Chu, B. R. Greene and G. Shiu, Mod. Phys. Lett. **A16**, 2231 (2001) [arXiv:hep-th/0011241].

[32] F. Lizzi, G. Mangano, G. Miele and M. Peloso, JHEP **0206**, 049 (2002) [arXiv:hep-th/0203099].

[33] R. H. Brandenberger, P. M. Ho, Phys. Rev. **D66**, 023517 (2002) [arXiv:hep-th/0203119].

[34] Q.-G. Huang and M. Li, JHEP **0306**(2003)014, hep-th/0304203; Q.-G. Huang and M. Li, JCAP **0311**(2003)001, astro-ph/0308458; Q. G. Huang and M. Li, Nucl. Phys. **B713**, 219 (2005) [arXiv:astro-ph/0311378].

[35] S. Tsujikawa, R. Maartens and R. Brandenberger, Phys. Lett. **B574**, 141 (2003) [arXiv:astro-ph/0308169]; S. Koh and R. H. Brandenberger, arXiv:hep-th/0702217; R. H. Brandenberger, arXiv:hep-th/0703173.

[36] R.-G. Cai, Phys. Lett. **B593**, 1 (2004), hep-th/0403134; Q. G. Huang and M. Li, Nucl. Phys. **B755**, 286 (2006) [arXiv:astro-ph/0603782]; X. Zhang and F. Q. Wu, Phys. Lett. **B638**, 396 (2006) [arXiv:astro-ph/0604195]; Qing-Guo Huang, Phys. Rev. **D74**, 063513 (2006), astro-ph/0605442; X. Zhang, JCAP **0612**, 002 (2006) [arXiv:hep-th/0608207]; Q. G. Huang, JCAP **0611**, 004 (2006) [arXiv:astro-ph/0610389].

[37] H. S. Kim, G. S. Lee and Y. S. Myung, Mod. Phys. Lett. **A20**, 271 (2005) [arXiv:hep-th/0402018].

[38] G. Calcagni, Phys. Rev. **D70**, 103525 (2004) [arXiv:hep-th/0406006]; G. Calcagni, Phys. Lett. **B606**, 177 (2005) [hep-ph/0406057]; G. Calcagni and S. Tsujikawa, Phys. Rev. **D70**, 103514 (2004) [astro-ph/0407543].

[39] Y.-F. Cai and Y.-S. Piao, arXiv:gr-qc/0701114.

[40] W. Xue, B. Chen and Y. Wang, arXiv:0706.1843 [hep-th].

[41] A. Linde, arXiv:0705.0164 [hep-th].

[42] J. Martin and R. H. Brandenberger, Phys. Rev. **D63**, 123501 (2001) [arXiv:hep-th/0005209].